

Unit _____	Standardized Pressure Injury Prevention Protocol Checklist (SPIPP- Adult) 2.0	Date _____
ITEM	Completed Yes/No	COMMENT
<b>Assess risk factors for pressure injury to guide risk-based prevention</b>		
Significant current or anticipated mobility problems		
Use a structured risk assessment approach (e.g., Braden or other validated risk tool) on admission		
Reassess risk q shift and with significant change in condition		
Patient/family informed of PI risk and prevention plan		
Additional risk factors considered: Previous PI __, Localized pain __, Diabetes __, Poor perfusion __, Vasopressors __, Oxygenation deficits __, Increased Temp __, Advanced Age __, Spinal cord injury __, Neuropathy __, Surgery/procedure duration > 2 hrs. __, Critical illness __, Organ Failure __, Sepsis __, Mechanical vent __, Medical devices __, Sedation __		
<b>Assess Skin/Tissue for signs of skin damage and pressure injury</b>		
Assess skin (comprehensive, visual, palpation) upon admission and q shift for erythema, discoloration, edema, and temperature		Location(s):
Assess skin under medical devices q shift		Device(s):
Inspect heels q shift		
In people of color: Ensure adequate lighting and moisten/moisturize skin to augment visual		
Consider enhanced skin assessment methods- thermography, SEM, skin color chart		
<b>Preventative Skin Care- Manage moisture/Incontinence</b>		
Cleanse and apply appropriate moisture barriers promptly after each incontinent episode		
Avoid use of alkaline soaps/cleaners		
Consider urinary/fecal management systems for high-risk persons		
Single layer, breathable, high absorbency pads for incontinence		
Consider using low friction textiles		
Apply wicking material to skin folds when appropriate		
<b>Redistribute Pressure</b>		
Turn/reposition q 2-3 hours persons who do not have independent bed mobility and as required by individual needs and risk, unless contraindicated (Braden Activity/Mobility score 1 or 2)		
Use high specification reactive foam or reactive air mattress/overlay for immobile persons (Braden Activity/Mobility score 1 or 2)		
Use positioning aids that minimize friction/shear (pillows, wedges). Use turn/lift equipment if available		
Keep head of bed as flat as possible		
Place silicone multilayer foam dressings on areas of high-risk (i.e., sacrum, lower buttocks, or heels) (Braden Activity/Mobility scores 1-2)		
Elevate heels off bed with pillows, heel devices or boots (Braden Sensory Perception score 1-3)		
Provide adequate repositioning (30 degree) when side lying		
Use slow, gradual, frequent, small, body shifts when unstable		
Use pressure redistributing seat cushion for persons who cannot adequately reposition		
Reposition seated persons q 1 hour		
Consult Physical Therapy for mobilization program when appropriate (Braden Activity/Mobility		
Consider reminder systems, pressure mapping, motion sensors		
Implement early mobilization program		
<b>Nutrition</b>		
Screen for malnutrition using a validated tool on admission		
Consult dietitian for persons with or at risk of malnutrition, decreased nutrient intake, NPO > 48 hours or presence of stage 2 or greater PI (Braden Nutrition Score 1-2)		
Provide additional calories, protein, fluids, and additional nutrients (i.e. multi-vitamin, arginine, glutamine, HMB) per nutrition plan of care or as appropriate		
Continue to regularly assess goals and consult dietitian as needed		