

## MG composite scale

Ptosis, upward ease (physician examination)	>45 seconds = 0	<b>11-45 seconds = 1</b>	1-10 seconds = 2	Immediate = 3
Double vision on lateral Gaze, left or right (physician examination)	>45 seconds = 0	11-45 seconds = 1	<b>1-10 seconds = 3</b>	Immediate = 4
Eye closure (physician examination)	Normal = 0	Mild weakness (can be forced open with effort) = 0	<b>Moderate weakness (can be forced open easily) = 1</b>	Severe weakness (unable to keep eye closed) = 2
Talking (patient history)	<b>Normal = 0</b>	Intermittent slurring or nasal speech = 2	Constant slurring or nasal but can be understood = 4	Difficult to understand speech = 6
Chewing (patient history)	<b>Normal = 0</b>	Fatigue with solid food = 2	Fatigue with soft food = 4	Gastric tube = 6
Swallowing (patient history)	Normal = 0	<b>Rare episode of choking or trouble swallowing = 2</b>	Frequent trouble swallowing, for example necessitating change in diet = 5	Gastric tube = 6
Breathing (thought to be caused by MG)	Normal = 0	<b>Shortness of breath with exertion = 2</b>	Shortness of breath at rest = 4	Ventilator dependence = 9
Neck flexion or extension (weakest) (physician examination)	Normal = 0	Mild weakness = 1	<b>Moderate weakness (i.e., ~50% weak, ±15%) = 3</b>	Severe weakness = 4
Shoulder abduction (physician examination)	Normal = 0	Mild weakness = 2	<b>Moderate weakness (i.e., ~50% weak, ±15%) = 4</b>	Severe weakness = 5
Hip flexion (physician examination)	Normal = 0	Mild weakness = 2	<b>Moderate weakness (i.e., ~50% weak, ±15%) = 4</b>	Severe weakness = 5
<b>TOTAL</b>				<u>20</u>

Note: Please note that “moderate weakness” for neck and limb items should be construed as weakness that equals roughly 50% ± 15% of expected normal strength. Any weakness milder than that would be “mild,” and any weakness more severe than that would be classified as “severe.”